

KANAZAWA-01-03

April, 2001

# The origin of quark and lepton mixings

Daijiro Suematsu \*

*\*Institute for Theoretical Physics, Kanazawa University,  
Kanazawa 920-1192, JAPAN*

## Abstract

We propose a model for the mass matrices of quarks and leptons based on two Abelian flavor symmetries. One is assumed to be broken at a high energy region near the Planck scale. It is used for the Froggatt-Nielsen mechanism in both quark and charged lepton sectors. Another one remains unbroken to a multi-TeV region. The mixings among neutrinos and gauginos including the one of the new Abelian symmetry generate non-zero masses and mixings among neutrinos. A bi-maximal scheme for the neutrino oscillation can be realized together with suitable masses and CKM-mixings in the quark sector. FCNC constraints on this flavor dependent Abelian symmetry seems to be evaded.

---

\*e-mail:suematsu@hep.s.kanazawa-u.ac.jp

# 1. Introduction

The origin of flavor mixings of quarks and leptons is one of the most important problems beyond the standard model (SM). Recently the existence of non-trivial lepton mixings has been strongly suggested through the atmospheric and solar neutrino observations whose results can be explained by assuming the neutrino oscillations [1, 2, 3]. The predicted flavor mixing is much bigger than the one of the quark sector. The explanation of this feature is a challenge to the grand unified theory (GUT) and a lot of works have been done by now [4, 5]. In many models the flavor mixing in both sectors is considered to be controlled by the Froggatt-Nielsen mechanism [6]<sup>1</sup> and the smallness of the neutrino mass is explained by the celebrated seesaw mechanism [8]. In such scenarios the origin of the flavor mixings is eventually related to the physics at high energies such as the Planck scale and an intermediate scale.

In this paper we propose an alternative possibility based on a different origin of the flavor mixings of neutrinos in a supersymmetric model with an extended gauge structure. In our scenario the flavor mixings in the quark and charged lepton sectors are considered to have its origin at the high energy region due to the usual Froggatt-Nielsen mechanism. On the other hand, the flavor mixings in the neutrino sector are assumed to come from their mixings with the extended gaugino sector by an extra U(1) gauge symmetry [9]. The mixings are induced by the R-parity violation [10, 11]. Its origin might be considered to be related to the physics concerning to an effective supersymmetry breaking at a TeV region. One of the interesting points of the model is that the large mixing angle MSW solution for the solar neutrino problem can be consistently accommodated together with the small flavor mixings and the qualitatively favorable mass eigenvalues in the quark sector. The scenario could be consistent with the gauge coupling unification since it has an SU(5) GUT structure when we switch off the low energy extra U(1) gauge interaction which plays a part of flavor symmetry together with the Froggatt-Nielsen type U(1)-symmetry. For convenience, we will use the SU(5) representations to classify the fields in the following discussion.

In the next section we define the flavor symmetry of the model. In section 3 the mass matrices in the quark and charged lepton sectors are discussed. FCNC constraints are

---

<sup>1</sup>There are many works in which the Abelian flavor symmetry is discussed. As examples, see [7] and references therein.

also examined for the non-universal couplings of an extra neutral gauge boson here. The neutrino mass matrix is studied in section 4. We show that a bi-maximal mixing is derived in this model. We also discuss the  $R$ -parity violation here. Section 5 is devoted to the summary.

## 2. Abelian flavor symmetry

We consider a model with two Abelian flavor symmetries  $U(1)_F \times U(1)_X$ . The  $U(1)_F$  is considered to be broken near the Planck scale and used to generate the mass hierarchy and the flavor mixing through the Froggatt-Nielsen mechanism. It may be considered to be an anomalous  $U(1)$  symmetry. On the other hand, the  $U(1)_X$  is assumed to remain unbroken to the TeV region<sup>2</sup>. We assume that this low energy extra  $U(1)_X$  gauge field has flavor diagonal but non-universal couplings. Different charges of  $U(1)_X$  are assigned to the  $\mathbf{5}^*$  fields belonging to the different generations<sup>3</sup>. Since its breaking scale is in the TeV region, we cannot use this symmetry for the Froggatt-Nielsen mechanism to induce the hierarchical structure of quark mass matrices since its breaking scale is too small as compared to the Planck scale. However, since a part of the contents of the minimal supersymmetric SM (MSSM) is assumed to have its charge in a generation dependent way, it can generate an additional non-trivial texture in the mass matrices.

We adopt the following charge assignments of  $U(1)_F \times U(1)_X$  to the chiral superfields of quarks and leptons:

$$\begin{aligned} \mathbf{10}_f &\equiv (q, u^c, e^c)_f & : & (3, 2, 0), \quad (\alpha, \alpha, \alpha), \\ \mathbf{5}^*_f &\equiv (d^c, \ell)_f & : & (1, 0, 0), \quad (q_1, q_1, q_2). \end{aligned} \tag{1}$$

where  $f=1-3$  and the numbers in the parentheses stand for the charges for each generation. We need no right-handed neutrino since in the present scenario neutrino masses are assumed to be generated through the mixings with gauginos. In general, the introduction of  $U(1)_X$  to the MSSM requires additional chiral superfields to cancel the gauge anomaly which causes the non-trivial constraint on the charge assignment. Since it is assumed to be radiatively broken at the TeV region, we need at least a new SM singlet chiral superfield whose scalar component causes the spontaneous breaking of  $U(1)_X$  by its vacuum

---

<sup>2</sup>Additional  $U(1)$ -symmetries are known to appear very often in the heterotic superstring models and it can play some useful roles in the supersymmetric models [12].

<sup>3</sup>This kind of charge assignment of  $U(1)_X$  has often been discussed in the different context. For example, it has been assumed to explain the small neutrino mass and the proton stability in [13].

expectation value (VEV). Taking account of these, as the Higgs chiral superfield sector we consider the following contents:

$$\begin{aligned}
\mathbf{5}^a &\equiv (D; H_2)^a & : & ((0, 0); (0, 0)), \quad ((x, z); (p, r)), \\
\mathbf{5}^{*a} &\equiv (\bar{D}; H_1)^a & : & ((0, 0); (0, 0)), \quad ((y, w); (q, s)), \\
\mathbf{1}_0 &\equiv S_0 & : & (-1), \quad (0), \\
\mathbf{1}_i &\equiv S_i & : & (0, \dots, 0), \quad (Q_1, \dots, Q_6),
\end{aligned} \tag{2}$$

where  $a=1-2$  and  $i=1-6$ . From the charge assignment of  $U(1)_X$  for  $\mathbf{5}^a$  and  $\mathbf{5}^{*a}$  we find that the  $SU(5)$  symmetry is explicitly broken unless  $U(1)_X$  is switched off. Then the  $SU(5)$  has only a meaning as a classification group in the model. The choice of additional chiral superfields and their charges should guarantee the SM gauge coupling unification, the proton stability and the anomaly cancellation of  $U(1)_X$ . The coupling unification of the SM gauge group is expected to be satisfied as a result of the  $SU(5)$  structure<sup>4</sup>. Moreover, the  $U(1)_X$  can prohibit the couplings between quarks and extra colored fields  $D^a$  and  $\bar{D}^a$  and guarantees the proton longevity.

Due to other phenomenological requirements some conditions should be imposed on the  $U(1)_X$ . It should allow the presence of necessary Yukawa couplings to generate the masses of quarks and leptons. For this requirement we impose the conditions

$$2\alpha + p = 0, \quad \alpha + q_1 + q = 0, \quad \alpha + q_2 + s = 0, \tag{3}$$

and then we have the following Yukawa couplings:

$$\begin{aligned}
W_{\text{Yukawa}} &= \sum_{f, f'=1}^3 y_{ff'}^u u_f^c H_2^1 q_{f'} \\
&+ \sum_{f'=1}^3 \left\{ \sum_{f=1,2} \left( y_{ff'}^d d_f^c H_1^1 q_{f'} + y_{ff'}^e e_f^c H_1^1 \ell_{f'} \right) + y_{3f'}^d d_{3f'}^c H_3^2 q_{f'} + y_{3f'}^e e_{3f'}^c H_3^2 \ell_{f'} \right\}. \tag{4}
\end{aligned}$$

We also require the following conditions on the Higgs chiral superfield sector:

$$\begin{aligned}
p + q + Q_1 &= 0, \quad p + s + Q_2 = 0, \quad x + y + Q_3 = 0, \quad Q_1 + Q_2 + Q_3 = 0, \\
r + q + Q_4 &= 0, \quad r + s + Q_5 = 0, \quad z + w + Q_6 = 0, \quad Q_4 + Q_5 + Q_6 = 0.
\end{aligned} \tag{5}$$

---

<sup>4</sup> The coupling unification requires an additional couple of  $SU(2)_L \times U(1)_Y$  vector like fields such as  $H_1$  and  $H_2$ . However, since they can be trivial under the  $U(1)_X$  and obtain masses, for example, due to the Giudice and Masiero mechanism [14], they can be irrelevant to our discussion.

We assume that the scalar components of the SM singlet fields  $S_3$  and  $S_6$  get VEVs at the TeV region radiatively through the couplings with extra colored fields  $D^a$  and  $\bar{D}^a$  and then the  $U(1)_X$  gauge field becomes massive [12]. Moreover, the last conditions in each line of (5) allow the trilinear couplings  $S_1 S_2 S_3$  and  $S_4 S_5 S_6$  in the superpotential. These trilinear couplings are accompanied with scalar trilinear couplings which break the supersymmetry softly and they can generally induce the VEVs to other singlet fields  $S_i$ . As a result all of the extra colored fields  $D^a$  and  $\bar{D}^a$  and the doublet Higgs fields become massive at that scale through the couplings with  $S_i$ . The mixings of the doublet Higgs fields in the superpotential can be written as

$$(H_2^1, H_2^2) \begin{pmatrix} \kappa_1 \langle S_1 \rangle & \kappa_2 \langle S_2 \rangle \\ \kappa_4 \langle S_4 \rangle & \kappa_5 \langle S_5 \rangle \end{pmatrix} \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix}. \quad (6)$$

If  $\frac{\kappa_5 \langle S_5 \rangle}{\kappa_4 \langle S_4 \rangle}$  is equal to  $-\frac{\kappa_1 \langle S_1 \rangle}{\kappa_2 \langle S_2 \rangle}$ , the eigenstates of the mixing matrix (6) can be identified as  $(H_2^1, H_1^\ell \equiv \sin \zeta H_1^1 + \cos \zeta H_1^2)$  and  $(H_2^2, H_1^h \equiv -\cos \zeta H_1^1 + \sin \zeta H_1^2)$  where  $\tan \zeta = \frac{\kappa_1 \langle S_1 \rangle}{\kappa_2 \langle S_2 \rangle}$ . Since  $H_2^1$  has a coupling with top quark as shown in eq. (4) and a mixing with  $H_1^\ell$  through eq. (6), only the former set of Higgs fields is expected to get the VEVs and it works as the usual Higgs fields. In the following discussion we take it as a basic assumption since the mixing in the  $H_1^a$  sector can play an important role to derive the MNS matrix due to the effect on the charged lepton mass matrix. Anomaly free conditions for  $U(1)_X$  can impose an additional constraint. If we require  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$  to be anomaly free under the conditions (3) and (5), the  $U(1)_X$  charges  $q_1$  and  $q_2$  will be constrained into the restricted region. The details are discussed in an appendix and we show a few examples of the solutions in Fig. 1.

### 3. Mass and mixing of quarks and charged leptons

The  $U(1)_F$  symmetry controls the flavor mixing structure by regulating the number of field  $S_0$  contained in each non-renormalizable term through the so-called Froggatt-Nielsen mechanism. As its result the effective Yukawa couplings  $y_{ff'}$  in eq. (4) have the hierarchical structure. In fact, if the singlet field  $S_0$  gets the VEV  $\langle S_0 \rangle$ , the suppression factor for the Yukawa couplings could appear as the power of  $\lambda = \frac{\langle S_0 \rangle}{M_{\text{pl}}}$ . Here  $M_{\text{pl}}$  is the Planck scale. Using the  $U(1)_F$  charges introduced above, we can obtain the mass matrices of quarks and charged leptons in the usual way. However, there is additional structure coming from the  $U(1)_X$  constraints which are realized by the condition (3) and

a composition of the doublet Higgs field  $H_1^\ell$ . Taking them into account we can write the mass matrices for the quarks and the charged leptons as follows:

$$M_u \sim \begin{pmatrix} \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^5 & \lambda^4 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix} \langle H_2^1 \rangle, \quad M_d \sim \begin{pmatrix} \lambda^4 \sin \zeta & \lambda^3 \sin \zeta & \lambda \sin \zeta \\ \lambda^3 \sin \zeta & \lambda^2 \sin \zeta & \sin \zeta \\ \lambda^3 \cos \zeta & \lambda^2 \cos \zeta & \cos \zeta \end{pmatrix} \langle H_1^\ell \rangle, \quad (7)$$

where the above mass matrices are written in the basis of  $\bar{\psi}_R m_D \psi_L$ . We do not consider the CP phases here. In eq. (7) we abbreviate the order one coupling constants by using the similarity symbol. In the quark sector the mass eigenvalues and the elements of the CKM matrix can be found after some inspection as

$$\begin{aligned} m_u : m_c : m_t &= \lambda^6 : \lambda^4 : 1, \\ m_d : m_s : m_b &= \lambda^4 \sin \zeta : \lambda^2 \cos \zeta : \cos \zeta, \\ V_{us} &\sim \lambda, \quad V_{ub} \sim \lambda^3, \quad V_{cb} \sim \lambda^2. \end{aligned} \quad (8)$$

If we require that the order one couplings should be in a range  $(\sqrt{\lambda}, \frac{1}{\sqrt{\lambda}})$ , the lower bound of  $\tan \beta \equiv \langle H_2^1 \rangle / \langle H_1^\ell \rangle$  is estimated as  $\tan \beta \gtrsim 40 \lambda \cos \zeta$ . On the charged lepton sector we can know the mass eigenvalues by noting that the SU(5) relation such as  $M_e^T = M_d$  is also satisfied in the present model. The ratio of mass eigenvalues is the same as the one of the down quark sector and then

$$m_e : m_\mu : m_\tau = \lambda^4 \sin \zeta : \lambda^2 \cos \zeta : \cos \zeta. \quad (9)$$

The result has some different features from the ones presented in ref. [5] in the down quark and charged lepton sectors. It comes from the charge assignment for  $\mathbf{5}^*$  and the composition of the Higgs field  $H_1^\ell$ . If we assume  $\lambda \sim 0.22$ , these results seem to describe nicely the experimental data for the mass eigenvalues and the CKM-mixing angles as far as  $\cos \zeta \sim \sin \zeta$  is satisfied, except for  $m_u$  and  $m_e$  which are predicted to be somehow larger. Also in our framework we cannot overcome this common defect with the scheme based on a kind of  $U(1)_F$  symmetry. The value of  $\sin \zeta$  is, in principle, determined by the scalar potential of the singlet fields  $S_i$  which is briefly discussed below (5). From its structure the above value of  $\sin \zeta$  seems to be expected to be realized without unnatural tunings of parameters. We define the diagonalization matrix  $\tilde{U}$  of the charged lepton mass

matrix in a basis that  $\tilde{U}^\dagger M_e^\dagger M_e \tilde{U}$  is diagonal. Then  $\tilde{U}$  can be approximately written as

$$\tilde{U} = \begin{pmatrix} 1 & 0 & \lambda \sin \zeta \\ -\lambda \sin^2 \zeta & \cos \zeta & \sin \zeta \\ -\lambda \sin \zeta \cos \zeta & -\sin \zeta & \cos \zeta \end{pmatrix}, \quad (10)$$

where the CP phase in the charged lepton sector is assumed to be zero although there can be some sources for it.

The breaking scale of  $U(1)_X$  is assumed to be in the TeV region and then we have a rather light  $Z'$  which can impose constraints on the model. The non-universal couplings of  $Z'$  with  $\mathbf{5}^*$  may induce large flavor changing neutral currents (FCNC). The detailed analysis for such an issue has been done in [15] and we can apply the discussion to the present model. In the model the  $Z'$  interaction term relevant to the non-universal couplings of  $\mathbf{5}^*$  can be written in the mass eigenstates as,

$$\begin{aligned} \mathcal{L}_{Z'} &= -g_1 \left[ \frac{g_X}{g_1} \cos \xi J_{(2)}^\mu - \sin \xi J_{(1)}^\mu \right] Z_\mu^{(2)}, \\ J_{(2)}^\mu &= \sum_{ij} \left[ \bar{\nu}_{L_i} B_{ij}^{\nu_L} \nu_{L_j} + \bar{\ell}_{L_i} B_{ij}^{\ell_L} \ell_{L_j} + \bar{d}_{R_i} B_{ij}^{\ell_R} d_{R_j} \right], \\ B_{ij}^\psi &= V^{\psi\dagger} \text{diag}(q_1, q_1, q_2) V^\psi, \end{aligned} \quad (11)$$

where  $J_{(1)}^\mu$  is the SM weak neutral current and  $\xi$  is a  $Z$ - $Z'$  mixing angle.  $V^\psi$  is a unitary matrix used to diagonalize the mass matrix of  $\psi$ . In the present model  $V^{d_R} = V^{\ell_L}$  is satisfied because of the  $SU(5)$  relation and they are represented by eq. (10). Thus the relevant  $B_{ij}^\psi$  can be estimated as

$$\begin{aligned} |B_{12}^{\ell_L}|^2 &= |B_{12}^{d_R}|^2 = (q_1 - q_2)^2 \lambda^2 \sin^4 \zeta \cos^2 \zeta, \\ |B_{13}^{\ell_L}|^2 &= |B_{13}^{d_R}|^2 = (q_1 - q_2)^2 \lambda^2 \cos^4 \zeta \sin^2 \zeta, \\ |B_{23}^{\ell_L}|^2 &= |B_{23}^{d_R}|^2 = (q_1 - q_2)^2 \cos^2 \zeta \sin^2 \zeta. \end{aligned} \quad (12)$$

On the other hand, the experimental constraints on these values are also estimated in [15]. The relevant constraints to the present model can be summarized as follows<sup>5</sup>. The coherent  $\mu$ - $e$  conversion and the decays  $\tau \rightarrow 3e, 3\mu$  require

$$w^2 (|B_{12}^{\ell_L}|^2) < 4 \times 10^{-14}, \quad w^2 (|B_{13}^{\ell_L}|^2) < 2 \times 10^{-5}, \quad w^2 (|B_{23}^{\ell_L}|^2) < 10^{-5}, \quad (13)$$

---

<sup>5</sup>We do not consider the CP violating effect here.

and from the lepton flavor violating meson decays such as  $K_L \rightarrow \mu^\pm e^\mp$  we obtain

$$y^2 |B_{12}^{\ell_L}|^2 |B_{12}^{d_R}|^2 < 10^{-14}. \quad (14)$$

The lepton flavor conserving meson decays such as  $K_L \rightarrow \mu^+ \mu^-$  impose

$$w^2 |\text{Re} [B_{12}^{d_R}]|^2 < 3 \times 10^{-11}, \quad (15)$$

and decays of  $B^0$  into  $\mu^+ \mu^-$  and  $\pi^0 \mu^+ \mu^-$  give

$$w^2 |B_{13}^{d_R}|^2 < 10^{-5}, \quad w^2 |B_{23}^{d_R}|^2 < 3 \times 10^{-6}. \quad (16)$$

In addition, from the experimental results on meson mass splittings we know the conditions

$$y |\text{Re} [(B_{12}^{d_R})^2]| < 10^{-8}, \quad y |\text{Re} [(B_{13}^{d_R})^2]| < 6 \times 10^{-8}, \quad y |\text{Re} [(B_{23}^{d_R})^2]| < 2 \times 10^{-8} \quad (17)$$

should be satisfied. In these conditions we use the definition

$$w = \frac{g_X}{g_1} (\rho_1 - \rho_2) \sin \xi \cos \xi, \quad y = \left( \frac{g_X}{g_1} \right)^2 (\rho_1 \sin^2 \xi + \rho_2 \cos^2 \xi), \quad (18)$$

where  $\rho_i = \frac{M_W^2}{M_{Z_i}^2 \cos^2 \theta_W}$ . If we apply the above value of  $\lambda$  and  $\sin \zeta \sim \frac{1}{\sqrt{2}}$  to eq. (12), the most stringent constraints on  $w$  and  $y$  are obtained as<sup>6</sup>

$$w^2 (q_1 - q_2)^2 < 7 \times 10^{-12}, \quad y (q_1 - q_2)^2 < 7 \times 10^{-12}. \quad (19)$$

If  $|q_1 - q_2|$  takes the value of order one,  $w$  and  $y$  should be smaller than  $10^{-6}$ . They require  $\sin \xi < 10^{-6}$  and  $M_{Z_2} > 100$  TeV. The value of charges  $q_1$  and  $q_2$  will be discussed from the view point of the explanation of the solar and atmospheric neutrino problems in the next section.

## 4. Neutrino mass and mixing

On the neutrino mass and mixing we adopt the scenario proposed in [9]. Using the  $U(1)_X$  charges defined in (1), the coupling between the neutrinos and the  $U(1)_X$  gaugino is given by  $i\sqrt{2}g_X \sum_\alpha q_\alpha (\tilde{\nu}_\alpha^* \lambda_X \nu_\alpha - \bar{\lambda}_X \bar{\nu}_\alpha \tilde{\nu}_\alpha)$ . We do not consider the kinetic term mixing between the  $U(1)$ -gauginos [16]. If we take this effect into account, off-diagonal elements appear in the gaugino mass matrix. The  $U(1)_X$  can plays an crucial role in the generation

---

<sup>6</sup>In this discussion we assume that the  $U(1)_X$ -charge is normalized in the same way as  $U(1)_Y$ .

of non-zero neutrino masses due to the above mentioned interaction since  $\nu_\tau$  has a charge  $q_2$  different from other neutrinos  $\nu_e$  and  $\nu_\mu$  whose charges are defined by  $q_1$ . If sneutrinos get the VEV  $u$  due to the R-parity violation, the mixing among neutrinos and gauginos appear as  $\mathcal{L}_{\text{mass}} = -\frac{1}{2}(\mathcal{N}^T \mathcal{M} \mathcal{N} + \text{h.c.})$  and

$$\mathcal{M} = \begin{pmatrix} 0 & m^T \\ m & M \end{pmatrix}, \quad m = \begin{pmatrix} a_2 & a_1 & b \\ a_2 & a_1 & b \\ a_2 & a_1 & c \end{pmatrix}, \quad M = \begin{pmatrix} M_2 & 0 & 0 \\ 0 & M_1 & 0 \\ 0 & 0 & M_X \end{pmatrix}, \quad (20)$$

where  $\mathcal{N}^T = (\nu_\alpha, -i\lambda_{W_3}, i\lambda_Y, -i\lambda_X)$ . We use the definition such as  $a_\ell = \frac{g_\ell}{\sqrt{2}}u$ ,  $b = \sqrt{2}g_X q_1 u$  and  $c = \sqrt{2}g_X q_2 u$ . If we assume  $u$  is much smaller than the gaugino masses  $M_A$ , we can obtain the light neutrino mass matrix from it by using the generalized seesaw formula. It can be written as

$$M_\nu = m^T M^{-1} m = \begin{pmatrix} m_0 + \epsilon^2 & m_0 + \epsilon^2 & m_0 + \epsilon\delta \\ m_0 + \epsilon^2 & m_0 + \epsilon^2 & m_0 + \epsilon\delta \\ m_0 + \epsilon\delta & m_0 + \epsilon\delta & m_0 + \delta^2 \end{pmatrix}, \quad (21)$$

where  $m_0, \epsilon$  and  $\delta$  are defined by

$$m_0 = \frac{g_2^2 u^2}{2M_2} + \frac{g_1^2 u^2}{2M_1}, \quad \epsilon = \frac{\sqrt{2}g_X q_1 u}{\sqrt{M_X}}, \quad \delta = \frac{\sqrt{2}g_X q_2 u}{\sqrt{M_X}}. \quad (22)$$

If the neutrinos have the same  $U(1)_X$  charge, we find that there is only one nonzero mass eigenvalue as in the usual R-parity violating scenario [10, 11]. The interesting aspect of this mass matrix is that it is defined only by the gaugino mass  $M_A$  ( $A = \ell, X$ ), the gauge couplings  $g_A$ , the  $U(1)_X$ -charges  $q_\alpha$  and the VEV  $u$  of sneutrinos. We define the mass eigenstates  $\nu_i$  by  $\nu_\alpha = U_{\alpha i} \nu_i$ . A mass eigenvalue  $m_2$  is zero and non-zero mass eigenvalues are represented as

$$m_{1,3} = \frac{1}{2} \left\{ (3m_0 + 2\epsilon^2 + \delta^2) \pm \sqrt{(m_0 + 2\epsilon^2 - \delta^2)^2 + 8(m_0 + \epsilon\delta)^2} \right\}. \quad (23)$$

Here we should note that the gaugino mass can have a CP phase which depends on the supersymmetry breaking mechanism. Because of it we can consider both possibilities of  $|m_1| < |m_3|$  and  $|m_1| > |m_3|$  depending on a sign of  $M_A$ . The diagonalization of the matrix (21) gives

$$U = \begin{pmatrix} \frac{1}{\sqrt{2}} \cos \theta & -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \sin \theta \\ \frac{1}{\sqrt{2}} \cos \theta & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \sin \theta \\ \sin \theta & 0 & \cos \theta \end{pmatrix}, \quad (24)$$

where one of the mixing angles  $\sin \theta$  is defined as

$$\sin^2 2\theta = \frac{8(m_0 + \epsilon\delta)^2}{(m_0 + 2\epsilon^2 - \delta^2)^2 + 8(m_0 + \epsilon\delta)^2}. \quad (25)$$

If we remind that the MNS-matrix which controls the neutrino oscillation phenomena is defined as  $V^{(\text{MNS})} = U^T \tilde{U}$ , we find that it can be written by using eqs. (10) and (24) as

$$\begin{pmatrix} \frac{\cos \theta}{\sqrt{2}}(1 - \lambda \sin^2 \zeta) - \lambda \sin \theta \sin \zeta \cos \zeta & -\frac{1}{\sqrt{2}}(1 + \lambda \sin^2 \zeta) & -\frac{\sin \theta}{\sqrt{2}}(1 - \lambda \sin^2 \zeta) - \lambda \cos \theta \sin \zeta \cos \zeta \\ \frac{\cos \theta}{\sqrt{2}} \cos \zeta - \sin \theta \sin \zeta & \frac{\cos \zeta}{\sqrt{2}} & -\frac{\sin \theta}{\sqrt{2}} \cos \zeta - \cos \theta \sin \zeta \\ \frac{\cos \theta}{\sqrt{2}}(1 + \lambda) \sin \zeta + \sin \theta \cos \zeta & \frac{1}{\sqrt{2}}(1 - \lambda) \sin \zeta & -\frac{\sin \theta}{\sqrt{2}}(1 + \lambda) \sin \zeta + \cos \theta \cos \zeta \end{pmatrix}. \quad (26)$$

Now we study the oscillation phenomena in the present model. It is well-known that the transition probability due to the neutrino oscillation  $\nu_\alpha \rightarrow \nu_\beta$  after the flight length  $L$  is written by using the matrix elements of (26) as

$$\mathcal{P}_{\nu_\alpha \rightarrow \nu_\beta}(L) = \delta_{\alpha\beta} - 4 \sum_{i>j} V_{\alpha i}^{(\text{MNS})} V_{\beta i}^{(\text{MNS})} V_{\alpha j}^{(\text{MNS})} V_{\beta j}^{(\text{MNS})} \sin^2 \left( \frac{\Delta m_{ij}^2}{4E} L \right), \quad (27)$$

where  $\Delta m_{ij}^2 = |m_i^2 - m_j^2|$ . If we remember  $m_2 = 0$ , we find that  $\Delta m_{12}^2$  or  $\Delta m_{23}^2$  should be relevant to the atmospheric neutrino problem. Unless both  $|\cos \theta| \sim 1$  and  $|\sin \zeta| \sim |\cos \zeta|$  are not satisfied, the relevant amplitude  $-4 \sum_{j=1,3} V_{\mu 2}^{(\text{MNS})} V_{\tau 2}^{(\text{MNS})} V_{\mu j}^{(\text{MNS})} V_{\tau j}^{(\text{MNS})}$  to  $\nu_\mu \rightarrow \nu_\tau$  is too small to explain the atmospheric neutrino deficit. Thus we assume  $\cos \theta \sim 1$  and  $\sin \zeta \sim \cos \zeta \sim \frac{1}{\sqrt{2}}$  here<sup>7</sup>. Under the assumption we obtain

$$V^{(\text{MNS})} \sim \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & -\frac{\lambda}{2} \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{\sqrt{2}} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (28)$$

This MNS-matrix is just the one representing the bi-maximal mixing [17, 18, 19]. In our model the large mixing between  $\nu_e$  and  $\nu_\mu$  has an origin in the neutrino sector and is related to the value of  $\cos \theta$ . On the other hand, the large mixing between  $\nu_\mu$  and  $\nu_\tau$  comes from the charged lepton sector and is relevant to the value of  $\cos \zeta$ . This feature makes it possible to meet the MNS matrix with the best fit value of the large mixing angle MSW solution (LMA) to the Super-Kamiokande data of the solar neutrino. It is a different situation from other bi-maximal mixing models based on the Zee model [23].

---

<sup>7</sup>The former condition requires the restricted values for  $q_1$  and  $q_2$  as is seen below. The latter one seems to be naturally satisfied as commented before.

On the squared mass differences the atmospheric neutrino deficit should be relevant to both  $\Delta m_{12}^2$  and  $\Delta m_{23}^2$ . On the other hand, the solar neutrino deficit should be related to  $\nu_e \rightarrow \nu_\mu$  with  $\Delta m_{12}^2$  and  $\nu_e \rightarrow \nu_\tau$  with  $\Delta m_{12}^2$ . It means that the normal hierarchy  $|m_2| \lesssim |m_1| \ll |m_3|$  should be satisfied and then  $M_A$  is required to have a negative sign in eq. (20). The CHOOZ experiment [20] constrains a component  $V_{e3}^{(\text{MNS})}$  of the MNS-mixing matrix since the amplitude of the contribution to  $\nu_e \rightarrow \nu_x$  with  $\Delta m_{23}^2$  or  $\Delta m_{13}^2$  always contains it. The model escapes the constraint since  $|V_{e3}^{(\text{MNS})}|$  takes  $\frac{\lambda}{2}$  which satisfies its upper bound 0.16 [21]. The value of  $V_{e3}^{(\text{MNS})}$  is a prediction of our model. In the model the effective mass parameter which appears in a formula of the rate of neutrinoless double  $\beta$ -decay [22] is estimated as

$$|m_{ee}| = \left| \sum_j |V_{ej}^{\text{MNS}}|^2 e^{i\phi_j} m_j \right| \sim \frac{1}{2}|m_1| + \frac{\lambda^2}{4}|m_3|. \quad (29)$$

Since  $|m_1| = \sqrt{\Delta m_{\text{solar}}^2}$  and  $|m_3| = \sqrt{\Delta m_{\text{atm}}^2}$ , it seems to be difficult that  $m_{ee}$  will be within the reach in near future.

Whether the above values of  $\cos \theta$  and mass eigenvalues  $m_i$  can be consistently realized is a crucial point for the model. We can study it by using the parameters  $M_A$ ,  $g_A$ ,  $q_\alpha$  and  $u$ . In the usual soft supersymmetry breaking scenario the gaugino mass is considered to be universally produced as  $M_0$  at the unification scale. Its low energy value is determined by the renormalization group equations (RGEs). If we use the one-loop RGEs, the gaugino mass at a scale  $\mu$  can be expressed as

$$M_2(\mu) = \frac{M_0}{g_U^2} g_2^2(\mu), \quad M_1(\mu) = \frac{5}{3} \frac{M_0}{g_U^2} g_1^2(\mu), \quad (30)$$

where we assume the gauge coupling unification at the scale  $M_U$  and define a value of the gauge coupling at  $M_U$  as  $g_U$ . It is not unnatural to consider the gauge coupling of  $U(1)_X$  and its gaugino mass to be the same as the ones of  $U(1)_Y$ <sup>8</sup>. If we adopt this simplified possibility, we can find that  $m_{1,3}$  and  $\sin^2 2\theta$  can be written by using only  $q_\alpha$  and  $\frac{g_U^2}{|M_0|} u^2$  as

$$m_{1,3} = \frac{3}{5} \left( -(2 + 2q_1^2 + q_2^2) \pm \sqrt{(2 + 2q_1^2 + q_2^2)^2 - \frac{16}{3}(q_1 - q_2)^2} \right) \frac{g_U^2 u^2}{|M_0|},$$

$$\sin^2 2\theta = \frac{8(2 + 3q_1 q_2)^2}{(2 + 6q_1^2 - 3q_2^2)^2 + 8(2 + 3q_1 q_2)^2}, \quad (31)$$

---

<sup>8</sup>In the superstring context the freedom of an Abelian Kac-Moody level can make it possible.




Fig. 1 Scatter plots of the  $U(1)_X$ -charge of neutrinos which can give the solution to the atmospheric and solar neutrino problems. Solid, dashed and dot-dashed lines are the ones corresponding to  $x = 1.5, 3$  and  $4.5$ , respectively.

where we assume  $M_0 < 0$ . The structure of the mass spectrum and the flavor mixing is controlled by the  $U(1)_X$ -charge. The gaugino mass  $M_0$  and the VEV  $u$  of sneutrinos are relevant to the mass eigenvalues only in the form of an overall factor  $\frac{g_U^2}{|M_0|}u^2$ . In order to realize the value in the range of  $2 \times 10^{-3} \text{ eV}^2 \lesssim \Delta m_{13}^2 \simeq \Delta m_{23}^2 \lesssim 6 \times 10^{-3} \text{ eV}^2$  which is required by the atmospheric neutrino deficit,  $\frac{g_U^2}{|M_0|}u^2$  needs to be in the range of  $0.003 \text{ eV} \lesssim \frac{g_U^2}{|M_0|}u^2 \lesssim 0.014 \text{ eV}$ . If we take  $|M_0| \sim 100 \text{ TeV}$ <sup>9</sup> and  $g_U \sim 0.72$ , for example, it shows that the sneutrino VEV should be  $u \sim 0.76 - 1.6 \text{ MeV}$ . The remaining freedom which we can use to explain the solar neutrino deficit is the  $U(1)_X$ -charge of  $\mathbf{5}_f^*$ . In Fig. 1 we plot the value of the  $U(1)_X$ -charge to realize the LMA for the explanation of the solar neutrino deficit which requires  $\cos \theta \sim 1$  as discussed above. Here we require  $\cos \theta > 0.98$  and also  $0.1 \times 10^{-4} \text{ eV}^2 \lesssim \Delta m_{12}^2 \lesssim 6 \times 10^{-4} \text{ eV}^2$  to draw the figure. It shows that the reasonable value of the  $U(1)_X$ -charge can realize the LMA. We should also note that the  $U(1)_X$  charges obtained here can satisfy the FCNC constraint given in (19) coming from the non-universal couplings of  $Z'$  as far as  $\sin \xi < 10^{-6}$  and  $M_{Z_2} > 100 \text{ TeV}$  are satisfied.

One of the unsolved important points is an origin of the small VEV  $u$  of sneutrinos. As

---

<sup>9</sup> The gaugino mass is required to be larger than the usually considered value. This comes from the  $M_{Z_2}$  value imposed by the FCNC constraint which has been discussed in the previous section.

mentioned in the previous part, it should be around  $O(1)$  MeV which is much smaller than the weak scale. In the MSSM there are arguments on the lepton number violation due to the VEVs of sneutrinos in the vicinity of the weak scale [24] and also some authors point out that the neutrino mass produced by them can be sufficiently small [25]. However, in our scenario we need much smaller VEVs of sneutrinos than the weak scale. To consider such a possibility it is useful to note that the small VEVs of sneutrinos could be obtained if there are bi-linear  $R$ -parity violating terms  $\epsilon L_\alpha H_2^1$  with a sufficiently small  $\epsilon$ . For example, as such a candidate we may consider the non-renormalizable terms, which are consistent with the  $U(1)_F \times U(1)_X$  symmetry such as

$$\frac{\bar{N}N}{M_{\text{pl}}^2} \frac{S_0}{M_{\text{pl}}} S_{\epsilon_1} L_1 H_2^1, \quad \frac{\bar{N}N}{M_{\text{pl}}^2} S_{\epsilon_2} L_2 H_2^1, \quad \frac{\bar{N}N}{M_{\text{pl}}^2} S_{\epsilon_3} L_3 H_2^1, \quad (32)$$

where new SM singlet fields  $S_{\epsilon_\alpha}$  are introduced for the  $U(1)_X$ -invariance. If an intermediate scale is induced through the  $D$ - and  $F$ -flat direction  $\bar{N} = N$  of other singlet fields  $N$ ,  $\bar{N}$  [12] and  $S_{\epsilon_\alpha}$  acquires the VEV at the TeV scale, the appropriate  $\epsilon$  term might be obtained. However, their equality is not guaranteed in the present example. Once we find that  $\epsilon$  terms could exist, we can check that the small  $u$  is realized by minimizing the scalar potential. Under the assumption that  $\langle H_1^\ell \rangle$  and  $\langle H_2^1 \rangle$  can be treated as constants, the value of  $u$  derived from the potential minimization is approximately expressed as

$$u \sim \epsilon \frac{\mu \langle H_1^\ell \rangle + B_\epsilon \langle H_2^1 \rangle}{m^2}, \quad (33)$$

where  $B_\epsilon$  is a soft supersymmetry breaking parameter related to the  $\epsilon L_\alpha H_2^1$  terms and  $m^2$  is the soft scalar mass of sneutrinos, which is taken to be universal here. From this we find that the sufficiently small  $u$  can be obtained as far as  $\epsilon$  is small enough and the  $\mu$ -parameter,  $B_\epsilon$  and  $m^2$  take the values of the order of weak scale. We need to check whether these conditions are satisfied at the true vacuum by taking account of the radiative correction. Even if the VEVs of sneutrinos are not equal and eq. (9) is somehow modified, which may be expected in the case of (32), there are two non-zero mass eigenvalues and the similar result obtained above might be derived as far as the deviations from the equal  $u$  are not so large. The quantitative analysis on this aspect is also necessary to know the viability of the model. Supersymmetry breaking scenario is also important for the model. These issues are now under investigation.

## 5. Summary

We have proposed the scenario for the origin of the mixings of quarks and leptons in the supersymmetric model with an extra  $U(1)_X$ -symmetry. The scenario is based on the usual Froggatt-Nielsen mechanism for the quark and charged lepton sectors. On the other hand, the mixing of neutrinos is considered to come from the mixing among neutrinos and gauginos induced by the R-parity violation. In this model we could obtain two non-zero mass eigenvalues for neutrinos at the tree level. The atmospheric and solar neutrino deficits can be simultaneously explained by the usual mass hierarchy scenario. In particular, the large mixing angle MSW solution for the solar neutrino problem can be realized consistently with the small quark mixings only by tuning the  $U(1)_X$ -charge of neutrinos. One of the important feature of the model is the value of  $V_{e3}^{(MNS)}$  which can take a rather large value within the CHOOZ constraint. The scenario gives an alternative possibility to the flavor mixing of quarks and leptons as compared to the usual one. That is, although the quarks and charged leptons have the origin of their mixings at the high energy region, the neutrinos may have it at the low energy region. The most crucial unsolved problem is to clarify the R-parity violation and the realization of the small sneutrino VEVs quantitatively. Further investigation of the problems seems to be necessary to see whether our model works well in a realistic way.

## Acknowledgement

The Author thanks M. Tanimoto for his useful comments. This work is supported in part by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (No.11640267).

## Appendix

Non-trivial anomaly free conditions in the model are summarized as

$$\begin{aligned}
\text{SU}(3)^2\text{U}(1)_X &: 9\alpha + 2q_1 + q_2 + x + y + z + w = 0, \\
\text{SU}(2)^2\text{U}(1)_X &: 9\alpha + 2q_1 + q_2 + p + q + r + s = 0, \\
\text{U}(1)_Y^2\text{U}(1)_X &: 45\alpha + 5(2q_1 + q_2) + 2(x + y + z + w) + 3(p + q + r + s) = 0, \\
\text{U}(1)_Y\text{U}(1)_X^2 &: p^2 - q^2 + r^2 - s^2 + y^2 - x^2 + w^2 - z^2 = 0, \\
\text{U}(1)_X^3 &: 30\alpha^3 + 5(2q_1^3 + q_2^3) + 3(x^3 + y^3 + z^3 + w^3) + \\
&\quad + 2(p^3 + q^3 + r^3 + s^3) + \sum_{i=1}^6 Q_i^3 = 0.
\end{aligned} \tag{34}$$

Combining eqs. (34) except for the last one with eqs. (3) and (5), we can express other parameters by  $q_1$ ,  $q_2$  and  $x$  as follows,

$$\begin{aligned}
\alpha &= -\frac{1}{9}(2q_1 + q_2), & p &= \frac{2}{9}(2q_1 + q_2), \\
q &= \frac{1}{9}(-7q_1 + q_2), & r &= \frac{1}{9}(q_1 + 5q_2), & s &= \frac{1}{9}(2q_1 - 8q_2), \\
y &= -\frac{1}{3}(q_1 - q_2) - x, & z &= -\frac{1}{3}(q_1 - q_2) + x, & w &= \frac{2}{3}(q_1 - q_2) - x, \\
Q_1 &= \frac{1}{3}(q_1 - q_2), & Q_2 &= -\frac{2}{3}(q_1 - q_2), & Q_3 &= \frac{1}{3}(q_1 - q_2), \\
Q_4 &= \frac{2}{3}(q_1 - q_2), & Q_5 &= -\frac{1}{3}(q_1 - q_2), & Q_6 &= -\frac{1}{3}(q_1 - q_2).
\end{aligned} \tag{35}$$

If we use eqs. (35) in the last one of eqs. (34) to solve it numerically, we can plot the solutions in the  $(q_1, q_2)$  plane for each value of  $x$ . In Fig. 1 we show it for three values of  $x$ .

## References

- [1] Y. Fukuda *et al.*, Phys. Rev. Lett. **81** (1998) 1562; R. Becker-Szendy *et al.*, IMB collaboration, Nucl. Phys. **B38** (proc. suppl.) (1995) 331; W. W. M. Allison *et al.*, Soudan collaboration, Phys. Lett. **B391** (1997) 491; M. Ambrosio *et al.*, Phys. Lett. **B434** (1998) 451.
- [2] B. T. Cleveland *et al.*, Ap. J. **496** (1998) 505; K. S. Hirata *et al.*, Kamiokande collaboration, Phys. Rev. Lett. **77** (1996) 1683; W. Hampel *et al.*, GALLEX collaboration, Phys. Lett. **B388** (1996) 384; J. N. Abdurashitov *et al.*, SAGE collaboration, Phys. Rev. Lett. **77** (1996) 4708.
- [3] Y. Suzuki, Super-Kamiokande collaboration, Talk presented at *Neutrino 2000*, Sudbury, Canada.
- [4] S. Bludman, N. Hata, D. Kennedy and P. Langacker, Nucl. Phys. Proc. Supp. **31** (1993) 156; K. S. Babu and S. M. Barr, Phys. Lett. **B381** (1996) 202; J. Sato and T. Yanagida, Phys. Lett. **B430** (1998) 127; B. Brahmachari and R. N. Mohapatra, Phys. Rev. **D58** (1998) 015001; C. Albright, K. S. Babu, and S. Barr, Phys. Rev. Lett. **81** (1998) 1167; K. S. Babu, J. C. Pati and F. Wilczek, Nucl. Phys. **B566** (2000) 33; M. Bando, T. Kugo and K. Yoshioka, Phys. Rev. Lett. **80** (1998) 3004; K. Oda, E. Takasugi, M. Tanaka and M. Yoshimura, Phys. Rev. **D59** (1999) 055001; T. Blazek, S. Raby and K. Tobe, Phys. Rev. **D60** (1999) 113001; Q. Shafi and Z. Tavartkilidze, Phys. Lett. **B487** (2000) 145; K. S. Babu and S. M. Barr, Phys. Rev. Lett. **85** (2000) 1170.
- [5] G. Altarelli and F. Feruglio, Phys. Lett. **B451** (1999) 388.
- [6] C. Froggatt and H. B. Nielsen, Phys. Lett. **B147** (1979) 277.
- [7] P. P. Binétruy, S. Lavignac, S. Petcov and P. Ramond, Nucl. Phys. **B496** (1997) 3; P. Binétruy, N. Irges, S. Lavignac and P. Ramond, Phys. Lett. **B403** (1997) 38; Y. Grossman, Y. Nir and Y. Shadmi, JHEP **9810** (1998) 007; Y. Nir and Y. Shadmi, JHEP **9905** (1999) 023.
- [8] M. Gell-Mann, P. Romond and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979) p.315; T. Yanagida, in *Proc.*

*Workshop on Unified Theory and Baryon Number in the Universe*, eds. O. Sawada and A. Sugamoto (KEK, 1979).

- [9] D. Suematsu, hep-ph/0012321 (to be published in Phys. Lett. **B**).
- [10] F. de Compos, M. A. Garcá-Jareño, A. S. Joshipura, J. Rosiek and J. W. F. Valle, Nucl. Phys. **B451** (1995) 3; T. Banks, Y. Grossman, E. Nardi and Y. Nir, Phys. Rev. **D52** (1995) 5319; A. S. Joshipura and M. Nowakowshi, Phys. Rev. **D51** (1995) 2421.
- [11] M. A. Díaz, J. C. Romão and J. W. F. Valle, Nucl. Phys. **B524** (1998) 23; J. W. F. Valle, in *Pro. 6th International Symposium on Particles, Strings and Cosmology* (PASCOS 98), Boston 1998, p502 and references therein.
- [12] D. Suematsu and Y. Yamagishi, Int. J. Mod. Phys. **A10** (1995) 4521; M. Cvetič and P. Langacker, Phys. Rev. **D54** (1996), Mod. Phys. Lett. **A11** (1996) 1247; D. Suematsu, Phys. Rev. **D59** (1999) 055017.
- [13] E. Nardi, Phys. Rev. **D48** (1993) 3277; Phys. Rev. **D49** (1994) 4394; E. Nardi and T. Rizzo, Phys. Rev. **D50** (1994) 203; D. Suematsu, Prog. Theor. Phys. **96** (1996) 611.
- [14] G. F. Giudice and A. Masiero, Phys. Lett. **B206** (1988) 480.
- [15] P. Langacker and M. Plumacher, Phys. Rev. **D62** (2000) 013006.
- [16] D. Suematsu, Mod. Phys. Lett. **A12** (1997) 1709; Phys. Lett. **B416** (1998) 108; Phys. Rev. **D57** (1998) 1738.
- [17] V. Barger, S. Pakvasa, T. J. Weiler and K. Whisnant, Phys. Lett. **B437** (1998) 107; S. Davidson and S. F. King, Phys. Lett. **B445** (1998) 191; R. N. Mohapatra and S. Nussinov, Phys. Rev. **D60** (1999) 013002.
- [18] I. Stancu and D. V. Ahluwalia, Phys. Lett. **B460** (1999) 431.
- [19] For the recent models with the large mixing, see for example, J. M. Mira, E. Nardi, D. A. Restrepo and J. W. F. Valle, Phys. Lett. **492** (2000) 81; A. de Gouvea and J. W. F. Valle, Phys. Lett. **B501** (2001) 115; P. H. Chankowski, A. Ioannisian,

- S. Pokorski and J. W. F. Valle, hep-ph/0011150; Q. Shafi and Z. Tavartkiladze, hep-ph/0101350; S. F. King and M. Oliveira, hep-ph/0009287.
- [20] C. Bemporad, CHOOZ collaboration, Nucl. Phys. B (Proc. Suppl.) **77** (1999) 159.
  - [21] G. Fogli, E. Lisi, A. Marrone and G. Scioscia, Phys. Rev. **D59** (1999) 033001; S. Bilenky, G. Giunti and W. Grimus, hep-ph/9809368.
  - [22] S. M. Bilenky, C. Giunti, C. W. Kim and S. T. Petcov, Phys. Rev. **D54** (1996) 4432; S. M. Bilenky, C. Giunti, W. Grimus, B. Kayser and S. T. Petcov, Phys. Lett. **B465** (1999) 193; H. V. Klapdor-Kleingrothaus, H. Päs and A. Yu. Smirnov, hep-ph/0003219.
  - [23] A. Zee, Phys. Lett. **93B** (1980) 389; Phys. Lett. **161B** (1985) 141.
  - [24] A. Masiero and J. W. F. Valle, Phys. Lett. **B251** (1990) 273; J. C. Romão, C. A. Santos and J. W. F. Valle, Phys. Lett. **B288** (1992) 311; J. C. Romão, A. Ioannisian and J. W. F. Valle, Phys. Rev. **D55** (1997) 427.
  - [25] J. C. Romão, M. A. Díaz, M. Hirsch, W. Porod and J. W. F. Valle, Phys. Rev. **D61** (2000) 071703; M. Hirsch, M. A. Díaz, W. Porod, J. C. Romão and J. W. F. Valle, Phys. Rev. **D62** (2000) 113008.